

FORMING CLUSTERS OF GALAXIES AS THE ORIGIN OF UNIDENTIFIED GeV GAMMA-RAY SOURCES

TOMONORI TOTANI¹ AND TETSU KITAYAMA²

¹National Astronomical Observatory, Mitaka, Tokyo 181-8588, Japan (e-mail: totani@th.nao.ac.jp)

²Department of Physics, Tokyo Metropolitan University, Hachioji, Tokyo 192-0397, Japan (e-mail: tkita@phys.metro-u.ac.jp)

Accepted by the Astrophysical Journal (Received 2000 May 9; Accepted August 3)

ABSTRACT

Over half of GeV gamma-ray sources observed by the EGRET experiment have not yet been identified as known astronomical objects. There is an isotropic component of such unidentified sources, whose number is about 60 in the whole sky. Here we calculate the expected number of dynamically forming clusters of galaxies emitting gamma-rays by high energy electrons accelerated in the shock wave when they form, in the framework of the standard theory of structure formation. We find that a few tens of such forming clusters should be detectable by EGRET and hence a considerable fraction of the isotropic unidentified sources can be accounted for, if about 5% of the shock energy is going into electron acceleration. We argue that these clusters are very difficult to detect in x-ray or optical surveys compared with the conventional clusters, because of their extended angular size of about $\sim 1^\circ$. Hence they define a new population of “gamma-ray clusters”. If this hypothesis is true, the next generation gamma-ray telescopes such as GLAST will detect more than a few thousands of gamma-ray clusters. It would provide a new tracer of dynamically evolving structures in the universe, in contrast to the x-ray clusters as a tracer of hydrodynamically stabilized systems. We also derive the strength of magnetic field required for the extragalactic gamma-ray background by structure formation to extend up to 100 GeV as observed, that is about 10^{-5} of the shock-heated baryon energy density.

Subject headings: cosmology: theory—diffuse radiation—large-scale structure of the universe—galaxies: clusters: general—gamma rays: theory

1. INTRODUCTION

The deepest image of the universe in the high energy gamma-ray band beyond 0.1 GeV has been obtained by the EGRET experiment (Hartman et al. 1999). Identified sources of the third EGRET catalog include the Large Magellanic Cloud, five pulsars, and 66 active galactic nuclei (AGNs) of the blazar class. However, over half of the EGRET sources have not yet been identified as known astronomical objects, and their origin is one of the most interesting mysteries in astrophysics. The distribution of these unidentified sources can be interpreted as the sum of the Galactic component along the Galactic disc ($|b| \lesssim 40^\circ$) and another isotropic (i.e., likely extragalactic) component (Özel & Thomson 1996; see also Fig. 2 of Mukherjee et al. 1995). Several candidates have been proposed as the origin of the Galactic unidentified sources, including molecular clouds, supernova remnants, massive stars, and radio-quiet pulsars (see e.g., Gehrels & Michelson 1999 and references therein). However, almost no candidate has been proposed to explain the extragalactic unidentified sources, except for undetected AGNs. Recently, Mirabal et al. (2000) have performed comprehensive follow-up observations for one of the high-latitude unidentified EGRET sources (3EG J1835+5918, $b = 25^\circ$) in X-ray, optical, and radio wavebands. They found that any known class of GeV gamma-ray sources including blazars and pulsars cannot

be the origin of 3EG J1835+5918, and it suggests that this source belongs to a new class of GeV gamma-ray emitters.

There are 19 unidentified sources with high galactic latitude of $|b| > 45^\circ$ (about 60 in the whole sky), and 7 of them are noted as ‘em’ in the third EGRET catalog, i.e., possibly extended or multiple sources that are inconsistent with single point sources. Although this ‘em’ designation is quite subjective and we should be careful in interpreting this result (see Hartman et al. 1999 for detail), this may indicate that there is an extended and extragalactic population in the unidentified EGRET sources. Recently Gehrels et al. (2000) chose ‘steady’ unidentified sources from the third EGRET catalog, to eliminate any source with high variability typical of flaring AGNs. These steady sources are mostly distributed at low and mid galactic latitude ($|b| \lesssim 40^\circ$) and hence should be the Galactic origin, but 7 sources are still located at $|b| > 45^\circ$, that may be steady extragalactic sources whose number is ~ 24 in the whole sky. Therefore, steady astronomical objects with an extended nature are worth being investigated as a possible origin of unidentified EGRET sources.

It is widely believed that the observed structures in the universe have been produced via gravitational instability. Currently the most successful theory of structure formation is the cold dark matter (CDM) scenario, in which the structures grow up hierarchically from small objects into larger ones. When an object collapses gravitationally

and virializes, the baryonic matter in the object is heated by shock waves up to the virial temperature. Particles are expected to be accelerated to high energy by shock acceleration, and accelerated electrons scatter the photons of the cosmic microwave background radiation (CMB) to high energy gamma-ray bands by the inverse-Compton mechanism. Existence of such nonthermal electrons is inferred from radio and hard x-ray observations for some clusters of galaxies (e.g., Fusco-Femiano et al. 1999), although the origin of the nonthermal electrons is not yet clear. It has also recently been argued that this radiation process in the intergalactic medium may explain the diffuse extragalactic gamma-ray background radiation (EGRB) observed in the EGRET range (Loeb & Waxman 2000). However, it is still highly speculative and difficult to test whether this process is really the origin of the EGRB, since the contribution by unresolved active galactic nuclei is also of the same order of magnitudes (see, e.g., Mücke & Pohl 2000 and references therein).

On the other hand, if the structure formation is actually an efficient radiation process of gamma-rays, clusters of galaxies should be strong emitters of gamma-rays when they dynamically form, and the detectability of such forming clusters as discrete sources is of great interest as a new probe of structure formation in the universe as well as a test for the scenario proposed by Loeb & Waxman (2000) for EGRB. In this paper we make a theoretical estimate of the number and angular size of such gamma-ray emitting clusters detectable by EGRET, based on the standard theory of structure formation in the CDM universe. We find a few tens of such forming clusters should have already been detected by EGRET. Detectability of such forming clusters in other wavebands such as optical or x-ray bands will be discussed, in comparison with the conventional clusters of galaxies identified in these wavebands. We will also calculate the EGRB spectrum from structure formation and derive a quantitative relation between the higher cut-off photon energy and magnetic field strength.

Throughout this paper, we assume a CDM universe with the density parameter $\Omega_0 = 0.3$, the cosmological constant $\Omega_\Lambda = 0.7$, the Hubble constant $h = H_0/(100\text{km/s/Mpc}) = 0.7$, the baryon density parameter $\Omega_B = 0.015h^{-2}$, and the density fluctuation amplitude $\sigma_8 = 1$. These cosmological parameters are consistent with various observations including those of the CMB fluctuations (e.g. de Bernardis et al. 2000) and the abundance of x-ray clusters of galaxies (e.g. Eke, Cole & Frenk 1996; Kitayama & Suto 1996b).

2. GAMMA-RAY LUMINOSITY AND FLUX FROM FORMING CLUSTERS

We first estimate the gamma-ray flux of a gravitationally bound object of total mass M that virializes at redshift z . The typical radius r_{vir} , density ρ_{vir} , circular velocity V_c , and temperature T_{vir} of the object can be computed from the spherical collapse model (Peebles 1980; Kitayama & Suto 1996b), that is widely used in study of structure formation. The total gravitational energy

given to the baryon gas in the forming cluster is given by $E_{\text{baryon}} \sim (3/4)(\Omega_B/\Omega_0)MV_c^2$. It is reasonable to expect that a fraction $\xi_e \sim 0.05$ of this energy goes into accelerated electrons, since such a fraction is inferred for acceleration of cosmic ray electrons in a supernova remnant SN 1006 from x-ray and TeV observations (Koyama et al. 1995; Tanimori et al. 1998) and consistent with the energetics among cosmic-rays, turbulent motions, and supernova rate in our Galaxy. It has also been suggested that the diffuse radio and hard x-ray emissions observed in the Coma cluster (and possibly other several clusters) can be attributed to nonthermal electrons with the electron energy fraction of the same order (e.g., Fusco-Femiano et al. 1999). Therefore, we use $\xi_e = 0.05$ to determine the normalization of electron energy spectrum throughout this paper.

The maximum Lorentz factor of electrons is constrained by the competition of the Fermi acceleration time and cooling time by inverse-Compton (IC) scattering of CMB photons. The acceleration time is given by

$$t_{\text{acc}} \sim \frac{r_L c}{V_s^2} \quad (1)$$

$$= 1.6 \times 10^{-4} \gamma_e B_{\mu G}^{-1} V_{s,3}^{-2} \text{ yr}, \quad (2)$$

where $r_L = m_e \gamma_e / (eB)$ is the Larmor radius of electrons, γ_e the electron Lorentz factor, e the electron charge, $B_{\mu G} = B/(1\mu G)$ the magnetic field, and $V_{s,3} = V_s/(10^3 \text{ km/s})$ the shock velocity that is of the same order of magnitudes with the circular velocity of a halo, V_c . On the other hand, the IC cooling time is

$$t_{\text{IC}} = \frac{\gamma_e m_e c^2}{(4/3) c \sigma_T U_{\text{CMB}} \gamma_e^2} \quad (3)$$

$$= 2.3 \times 10^{12} \gamma_e^{-1} (1+z)^{-4} \text{ yr}, \quad (4)$$

where σ_T is the Thomson cross section and $U_{\text{CMB}} = 4.32 \times 10^{-13} (1+z)^4 \text{ erg cm}^{-3}$ is the CMB energy density. Equating these expressions of t_{acc} and t_{IC} , we have the maximum value of γ_e as $\gamma_{e,\text{max}} = 1.2 \times 10^8 (1+z)^{-2} B_{\mu G}^{1/2} V_{s,3}$. We assume the energy distribution of accelerated electrons as a power-law with an exponential cut-off at $\gamma_{e,\text{max}}$, i.e., $dN_e/d\gamma_e \propto \gamma_e^{-\alpha} \exp(-\gamma_e/\gamma_{e,\text{max}})$, with the standard particle acceleration index of $\alpha \sim 2$. As mentioned above, the normalization of this spectrum is set by the equation

$$\int d\gamma_e m_e c^2 \gamma_e \frac{dN_e}{d\gamma_e} = \xi_e E_{\text{baryon}}, \quad (5)$$

with the parameter $\xi_e = 0.05$. The observed photon energy ϵ_γ scattered by electrons is related to γ_e as $\epsilon_\gamma = (4/3) \gamma_e^2 \epsilon_{\text{CMB},0}$, where $\epsilon_{\text{CMB},0} = 6.4 \times 10^{-4} \text{ eV}$ is the mean photon energy of the CMB at $z = 0$.

The cooling time of electrons corresponding to photon energy ϵ_γ can be written as $t_{\text{IC}} = 2.1 \times 10^6 (\epsilon_\gamma/\text{GeV})^{-1/2} (1+z)^{-4} \text{ yr}$, and this should be compared with the time for the shock wave to propagate the radius of the virialized halo, $t_{\text{shock}} \sim r_{\text{vir}}/V_s = r_{\text{vir}}/(4V_c/3)$. Here we have estimated the shock velocity as $V_s = (4/3)V_c$,

that is a velocity of a strong shock when a material is shocked by a supersonic piston with a velocity of V_c , i.e., a typical bulk velocity of material in a collapsed halo. By using the spherical collapse model mentioned above to calculate r_{vir} and V_c for the halo, this time scale can be written as $t_{\text{shock}} = (3/4)^{3/2} \pi^{-1/2} (G\rho_{\text{vir}})^{-1/2} \sim 1.5(1+z)^{-3/2}$ Gyr, that is essentially the dynamical time of the halo. Note that it depends only on the redshift and not on the halo mass. From this argument the cooling time of electrons emitting gamma-rays above 0.1 GeV is always much shorter than the time scale t_{shock} during which the shock is alive and a halo is an active gamma-ray emitter. Hence the total number of gamma-rays emitted from a forming halo during the time t_{shock} is given as:

$$\frac{dN_\gamma(\epsilon_\gamma; M, z)}{d\epsilon_\gamma} = \frac{m_e \gamma_e}{\epsilon_\gamma} \frac{dN_e}{d\gamma_e} \frac{d\gamma_e}{d\epsilon_\gamma}, \quad (6)$$

and the observed photon flux of gamma-rays during the shock propagation time is

$$\frac{dF(\epsilon_\gamma; M, z)}{d\epsilon_\gamma} = \frac{(1+z)}{4\pi d_L^2} \frac{dN_\gamma}{d\epsilon_\gamma} \frac{1}{t_{\text{shock}}}, \quad (7)$$

where d_L is the standard luminosity distance.

We introduce a parameter ξ_B to determine the magnetic field, which is the fraction of magnetic energy density in the total baryon energy density of the halo, i.e., $B^2/(8\pi) = \xi_B E_{\text{baryon}}/(4\pi r_{\text{vir}}^3/3)$. Then by using the spherical collapse model again, the magnetic field of a cluster can be written as

$$B \sim 0.17 \left(\frac{\xi_B}{10^{-3}} \right)^{1/2} \left(\frac{M}{10^{15} M_\odot} \right)^{1/3} (1+z)^2 \mu\text{G}. \quad (8)$$

A magnetic field of ~ 0.1 – $1 \mu\text{G}$ is often observed in intracluster medium of rich clusters (Kronberg 1994; Fusco-Femiano et al. 1999; Rephaeli, Gruber, & Blanco 1999), and hence we use $\xi_B = 10^{-3}$ to be consistent with the observations. It should be noted, however, that this parameter is important only for the maximum photon energy of the gamma-ray spectrum (well beyond 10–100 GeV with $\xi_B = 10^{-3}$, see §4), and the gamma-ray flux above 100 MeV measured by the EGRET is almost insensitive to this uncertain parameter, unless the particle index α significantly deviates from the standard value of 2.

It is interesting to apply the above model to nearby known clusters of galaxies. For example, the Coma cluster has the total mass of $M \sim 10^{15} M_\odot$ and is located at $z = 0.023$. The flux with these parameters becomes $F(> 0.1\text{GeV}) \sim 6.5 \times 10^{-7}$ photons $\text{cm}^{-2}\text{sec}^{-1}$, that is about 10 times brighter than the observational upper limit on this cluster by EGRET, 4×10^{-8} photons $\text{cm}^{-2}\text{sec}^{-1}$ (Sreekumar et al. 1996). This does not mean, however, that our model is incorrect. Our model is relevant for just-forming clusters of galaxies in which the violent shock generated by gravitational collapse is still alive. The Coma cluster is thought to have formed more than a few dynamical times ago and is now hydrodynamically stable after

violent shock has disappeared. Gamma-ray flux can then be much weaker than our estimation. On the other hand, it suggests that there was an epoch during which this cluster was a strong gamma-ray emitter, and also that there may be other clusters visible by EGRET, that are just dynamically forming and have not yet reached hydrostatic equilibrium.

3. EXPECTED NUMBER OF GAMMA-RAY CLUSTERS DETECTABLE BY EGRET

The number of such forming clusters of galaxies with flux stronger than F can be calculated as

$$N(> F) = \int dz \int_{M(z; F)}^{\infty} dM \frac{dV}{dz} R_{\text{form}}(M, z) t_{\text{shock}}, \quad (9)$$

where dV/dz is the comoving volume element of the universe, R_{form} the formation rate of dark haloes (or clusters) per unit mass, cosmic time, and comoving volume, and $M(z; F)$ the mass of a cluster collapsing at redshift z whose flux is F . Here we have taken into account that clusters are active gamma-ray emitters only during the time t_{shock} .

There are several formulae to calculate the formation rate R_{form} in the framework of the standard theory of structure formation. As is well known, the Press & Schechter (1974, PS; Peebles 1980) formalism provides a formula of mass function (i.e., number density of haloes as function of mass and redshift), that is in reasonable agreement with N -body simulations. Here we want the formation rate of haloes rather than the mass function at a given epoch, because we need to calculate the number of collapsing objects experiencing shock at each epoch. A naive prescription to obtain this quantity is to take a time derivative of the PS mass function, R_{dPS} , although this is not exactly R_{form} but rather interpreted as $R_{dPS} = R_{\text{form}} - R_{\text{dest}}$, where R_{dest} is the rate of destruction of haloes by merging into even larger structures. Consequently, R_{dPS} becomes negative at small mass scales where R_{dest} is significant. As shown below, however, the number of objects visible by EGRET is dominated by massive clusters forming in recent past ($z \ll 1$) that are the largest structures in the universe, and hence R_{dest} is negligible. Therefore it is a reasonable approximation to use R_{dPS} in the mass range where it is positive. Alternatively, there are several extensions to the PS theory for computing R_{form} (e.g., Lacey & Cole 1993; Sasaki 1994; Kitayama & Suto 1996a; Percival & Miller 1999). In what follows we use R_{dPS} and the formulae of Sasaki (1994) and Kitayama & Suto (1996a) to take account of theoretical uncertainties in R_{form} .

Figure 1 shows the theoretically predicted $\log N$ - $\log F$ of forming clusters. At least a few tens of clusters should be visible by the EGRET, and a significant fraction of the isotropic unidentified EGRET sources can be accounted for. It is also interesting to note that the predicted number is similar to that of ‘em’ isotropic unidentified EGRET sources, i.e., possibly extended sources. The predicted number at the EGRET flux limit also agrees with the num-




FIG. 1.— The upper panel: the cumulative flux distribution of gamma-ray emitting clusters of galaxies. The three curves are the theoretical predictions using three different formulae for the formation rate of dark haloes (R_{form}), i.e., the derivative of the Press-Schechter (PS) mass function (solid line), formation rate of Sasaki (1994; dot-dashed line) and Kitayama & Suto (1996a; dashed line). The dotted line is the flux distribution of uniformly distributed sources in the Euclidean space. The observed distribution of the unidentified EGRET sources with $|b| > 45^\circ$ is shown by the thick (only the ‘em’ sources) and thin (all sources including non-‘em’) lines. The lower panel: the solid, dashed, and dot-dashed lines are mean redshift, mass [$10^{16} M_\odot$], and angular radius [degree] of gamma-ray clusters brighter than a given flux, respectively. The dotted line is the contribution of gamma-ray clusters brighter than a given flux to the diffuse extragalactic gamma-ray background at 100 MeV. The derivative of the PS mass function is used. The sensitivity limits of the EGRET and GLAST experiments are shown in the figure.

ber of ‘steady’ unidentified sources with $|b| > 45^\circ$ defined by Gehrels et al. (2000). This result is robust against changes in the adopted prescription for R_{form} . In the lower panel of Fig. 1, we also show mean mass, redshift, and angular radius θ_{vir} corresponding to r_{vir} of such gamma-ray clusters brighter than a given flux. These quantities are $M \sim 10^{15} M_\odot$, $z \sim 0.05$, and $\theta_{\text{vir}} \sim 1^\circ$ for clusters above the EGRET sensitivity limit. Considering the EGRET angular resolution, the typical radius of $\sim 1^\circ$ is consistent with the fact that a significant fraction of isotropic unidentified sources are indicated as possibly extended.

It is predicted that more than a few thousands of forming clusters will be detected by future missions such as the GLAST (Gehrels & Michelson 1999), and the improved angular resolution may reveal the extended profile for nearby gamma-ray clusters with higher statistical significance. Another important prediction is that GLAST will observe the flattening of the $\log N$ - $\log F$ curve due to the cosmological effects, compared with the expectation of a uniform source distribution in the Euclidean space (dotted line in the upper panel of Fig. 1).

4. EXTRAGALACTIC GAMMA-RAY BACKGROUND

Our formulation also allows us to calculate the EGRB flux and spectrum as

$$\frac{dn_\gamma}{d\epsilon_\gamma} = \int dz \int dM \frac{dN_\gamma(\epsilon_\gamma; M, z)}{d\epsilon_\gamma} R_{\text{form}}(M, z) \frac{dt}{dz} \quad (10)$$

where $(dn_\gamma/d\epsilon_\gamma)$ is the gamma-ray number density that is related to the EGRB flux as $(dF/d\epsilon_\gamma) = c(dn_\gamma/d\epsilon_\gamma)/(4\pi)$, and t is the cosmic time. This flux becomes $\epsilon_\gamma^2(dF/d\epsilon_\gamma) \sim 1 \text{ keV cm}^{-2} \text{ sec}^{-1} \text{ sr}^{-1}$ at 100 MeV, in good agreement with the observation (Sreekumar et al. 1998) as well as the previous simpler estimation assuming that the average temperature of baryons in the universe is $\sim \text{keV}$ at present (Loeb & Waxman 2000). In fact, we have checked that the mass-averaged temperature of virialized haloes in the universe as a function redshift, that is calculated by the PS theory, agrees within a factor of 2 with a numerical simulation (Cen & Ostriker 1999) on which the previous EGRB estimate was based. This fact gives a justification for the use of the PS theory to calculate the gamma-ray emitting objects. The dotted line in the lower panel of




FIG. 2.— The spectrum of the diffuse extragalactic gamma-ray background radiation. The data is from Sreekumar et al. (1998). The parameter ξ_B is the fraction of magnetic energy in the gravitational energy given to baryonic gas in a collapsed halo, with $\xi_B = 10^{-1}$ (dashed line), 10^{-3} (solid), and 10^{-5} (dot-dashed). The absorption of gamma-rays above ~ 100 GeV by the pair-production interaction with the cosmic infrared background is not taken into account in the upper panel, while it is in the lower panel. (The secondary gamma-rays reprocessed by the produced pairs are not taken into account in either of the panels, see text.)

Fig. 1 shows the contribution to the EGRB at 100 MeV by objects brighter than a given flux. We predict that GLAST will resolve about 20–30 % of the EGRB as discrete gamma-ray clusters, if structure formation is the major origin of the EGRB.

The strength of magnetic field in the shocked baryons is important for the question whether the EGRB spectrum extends up to ~ 100 GeV as observed. In Fig. 2 we show the EGRB spectrum with several values of ξ_B . This result shows that the magnetic field strength corresponding to $\xi_B \sim 10^{-5}$ of the baryon energy density is sufficient for the EGRB to extend beyond 100 GeV. The magnetic field observed in the intracluster gas ($\xi_B \sim 10^{-3}$) is much stronger than this, and it is also theoretically reasonable to expect that the turbulent motion in collapsed objects amplifies the seed magnetic field made by the battery mechanism well beyond $\xi_B \gtrsim 10^{-5}$ within the dynamical time (Kulsrud et al. 1997). Therefore, physically reasonable magnetic field strength can explain the extension of the EGRB spectrum beyond 100 GeV, and it is likely that a considerable fraction of gamma-rays above 100 GeV is absorbed by the interaction with the cosmic infrared background radiation producing electron-positron pairs. The effect of intergalactic absorption is shown in

the lower panel of Fig 2, using the optical depth of intergalactic pair-production presented in Totani (2000). The absorbed TeV gamma-rays will be reprocessed into GeV gamma-rays by the produced pairs, and distort the EGRB spectrum (e.g., Coppi & Aharonian 1997). Although these secondary GeV gamma-rays are not taken into account here and it is beyond the scope of this paper, it is important to study how large is this spectral distortion in future work.

5. DISCUSSION

We here discuss the expected properties of gamma-ray clusters of galaxies. Perhaps the most natural question in this regard would be “Are they already observed in other wavebands such as x-rays or optical surveys?” We have checked that there is no statistically significant association of the ROSAT Brightest Cluster Sample (RBCS, Ebeling et al. 1998), within the 95% error circles of the unidentified sources with $|b| > 30^\circ$ in the EGRET catalog. We have also checked the correlation with the clusters in the revised Abell catalog (Abell, Corwin, & Olowin 1989), and no statistically significant associations are found, either. However, in the following we argue that the gamma-ray clusters proposed in this paper are very difficult to detect

in x-rays or optical bands compared with ordinary clusters identified in these wavebands, and hence our scenario is not rejected by these results.

5.1. Detectability of gamma-ray clusters in X-rays

We first estimate the expected x-ray flux from gamma-ray clusters. Baryonic gas in most clusters of galaxies observed in x-rays seems to be in approximate hydrostatic equilibrium with the surface brightness well fitted by a density profile, $\rho_{\text{gas}}(r) \propto [1 + (r/r_c)^2]^{-1}$ (e.g., Sarazin 1988), where r_c is the core radius that is typically about ~ 10 times smaller than the virial radius. Since the x-ray emissivity is proportional to ρ_{gas}^2 , the x-ray emission is strongly concentrated into the central region. Assuming the above density profile and the self-similar model as described in Kitayama & Suto (1996b),¹ a typical gamma-ray cluster detectable by the EGRET with $M \sim 10^{15} M_\odot$ and $z \sim 0.05$ would have the x-ray flux $2.4 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$ in 0.1–2.4 keV.

The inverse-Compton flux is also expected to be comparable with the thermal emission. By equating t_{IC} and t_{shock} in §2, we get the cooling photon energy $\epsilon_{\gamma, \text{cool}} = 2.0(1+z)^{-5} \text{ keV}$, below which the electron cooling time is longer than the dynamical time. Then the IC spectrum extends down to around x-ray band with $dN_\gamma/d\epsilon_\gamma \propto \epsilon_\gamma^{-2}$, while it becomes harder at wavelengths longer than x-rays with $dN_\gamma/d\epsilon_\gamma \propto \epsilon_\gamma^{-1.5}$. If the gamma-ray flux at 100 MeV is $\sim 10^{-7} \text{ photons cm}^{-2} \text{ s}^{-1}$ that is the EGRET threshold, the IC x-ray flux (νF_ν) is $\sim 1.6 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$. Therefore, the thermal and IC fluxes are well above the flux limit $\sim 4 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ of the RBCS.

However, it takes nearly the dynamical time for the cluster gas to reach hydrostatic equilibrium after the collapse, and gamma-rays from the shock generated by the gravitational collapse are radiated away within that period. Then it is likely that the density profile of gamma-ray emitting clusters is more irregular and extended than ordinary x-ray clusters. In fact, if the unidentified ‘em’ sources in the EGRET catalog are actually extended, they must have typical angular size of about degree, from the source location accuracy of the EGRET. As we have shown, angular size of about 1° is theoretically reasonable if the emission is extended to the virial radius. When the density profile is not concentrated into the central region but rather constant within the virial radius, the x-ray luminosity becomes lower than the self-similar model by a factor of ~ 3.7 because of the lower central density.

Furthermore, the surface brightness of such loose clusters should be drastically dimmer than ordinary x-ray clusters. In the self-similar model with $r_{\text{vir}} \gg r_c$, the core gas density is $\rho_{\text{gas}, c} \sim (1/3)(r_{\text{vir}}/r_c)^2 \rho_{\text{gas}, \text{vir}} \sim 50 \rho_{\text{gas}, \text{vir}}$, where $\rho_{\text{gas}, \text{vir}} = (\Omega_B/\Omega_0)\rho_{\text{vir}}$ is the virial gas density that is the average gas density within r_{vir} . On the other hand, if the gas density profile of gamma-ray clusters is roughly

constant at $\rho_{\text{gas}, \text{vir}}$ out to r_{vir} , the x-ray surface brightness of such a loose cluster is dimmer than the central surface brightness of the self-similar model by a factor of $\sim (r_c/r_{\text{vir}})(\rho_{\text{gas}, c}/\rho_{\text{gas}, \text{vir}})^2 \sim 200$, since the x-ray emissivity is proportional to ρ_{gas}^2 . It crucially affects the detectability of x-rays from gamma-ray clusters. The detectability of x-rays should be described by the signal-to-noise ratio (S/N) against the x-ray background flux. The noise level is proportional to $(\text{image area})^{1/2}$, and hence $S/N \propto F/r$, where F and r are the flux and the image radius, respectively. We have compared the value of F/r of the extended gamma-ray clusters detectable by the EGRET and those of the clusters in the RBCS. We found that the F/r of gamma-ray clusters is by a factor of 3 smaller than the minimum F/r of the RBCS clusters. The absence of association between the RBCS and the EGRET sources is therefore not in contradiction to our scenario. On the other hand, deeper observation of candidate gamma-ray clusters by Newton, for example, might detect the x-ray emission extended to about 1° with the flux estimated above, that would provide a clear test of our scenario. Such x-ray emission should reflect the structure of shocks in dynamically forming clusters, and imaging study is of great interest.

5.2. Detectability in the Optical Surveys

Here we again emphasize that the gamma-ray clusters are expected to be more extended than clusters that have already stabilized. It is known that the surface density profile of galaxies in a cluster can well be described by the King profile, $\sigma(r) \propto [1 + (r/r_c)^2]^{-1}$ with the core radius of $\sim 100 \text{ kpc}$ that is comparable with the core radius of x-ray profile (e.g., Adami et al. 1998). If we assume a roughly constant surface density out to $\sim r_{\text{vir}}$ rather than the King profile for gamma-ray clusters, the average surface density $\sigma_{\text{av}} \sim N_{\text{gal}}/(\pi r_{\text{vir}}^2)$ is lower than the central surface density of the King profile $\sigma_c \sim N_{\text{gal}}/[2\pi r_c^2 \ln(r_{\text{vir}}/r_c)]$, by a factor of $\sigma_c/\sigma_{\text{av}} \sim [2 \ln(r_{\text{vir}}/r_c)]^{-1}(r_{\text{vir}}/r_c)^2 \sim 30$. Here N_{gal} is the total number of galaxies within r_{vir} . This dimming factor is not so significant as that for the x-ray surface brightness, but that should make the optical identification very difficult because of the contamination by foreground and/or background field galaxies. Therefore, we consider that no statistically significant association with the known optically identified clusters does not immediately reject our scenario. Instead, it is necessary to study in the future the correlation between the EGRET sources and galaxy catalogs taking into account the possibility that the gamma-ray clusters are considerably extended. Search in optical bands has an advantage over the search in x-rays, in a sense that the dimming of surface number density compared with ordinary clusters is less severe than x-rays whose emissivity is proportional to ρ_{gas}^2 . The typical density of such loose clusters is close to the virial

¹Here we have assumed that the core radius is proportional to the virial radius as $r_c = 0.21(h/0.7)^{-1} \text{ Mpc} \times [r_{\text{vir}}(M, z)/r_{\text{vir}}(10^{15} M_\odot, 0)]$ with $r_{\text{vir}}(10^{15} M_\odot, 0) = 2.6 \text{ Mpc}$, where the normalization is chosen to match the local observations (Abramopoulos & Ku 1983; Jones & Forman 1984).

density, that is about a few hundreds times higher than the mean density of the universe.

We have also noticed that there are a considerable number of ‘em’ sources in the EGRET sources identified as AGNs. If they were actually extended sources, it might be speculated that some of them are also gamma-ray clusters including an AGN as a member galaxy. Time variability of these sources would be an important test to check this possibility.

5.3. *On the recent follow-up observations for 3EG J1835+5918*

Recent follow-up observations by Mirabal et al. (2000) for one of the high-latitude unidentified EGRET sources (3EG J1835+5918) have found a diffuse x-ray emission from an uncatalogued cluster of galaxies at $z = 0.102$. Although this cluster is outside the 99% error ellipse of 3EG J1835+5918 whose radius is $12'$, the separation between the centers of the x-ray cluster and 3EG J1835+5918 is about 0.65° , that is within our expectation of the typical angular radius of gamma-ray clusters detectable by the EGRET, $\sim 1^\circ$. As discussed above, x-ray emission is expected from a region where the intracluster gas reached hydrodynamical equilibrium, while gamma-rays are emitted from a region still hydrodynamically unstable. Therefore, it is not surprising that the positions of x-ray and gamma-ray emissions are different unless the separation is well beyond the virial radius of $\sim 1^\circ$.

3EG J1835+5918 is not an ‘em’ source, and this source may not be an extended source. The radius of the 99% confidence ellipse, 0.2° , is considerably smaller than the expected angular size of gamma-ray clusters. However, high energy electrons emitting GeV gamma-rays have very short life time compared with the shock propagation time ($t_{IC} \sim 10^{-3}t_{shock}$), and gamma-ray emitting region may be very clumpy in a cluster. (On the other hand, x-ray emitting electrons have a cooling time comparable with or longer than t_{shock} for IC and thermal radiations, and hence x-ray emitting region should be much less clumpy and extended with the size $\sim \theta_{vir}$, as discussed in §5.1.) Therefore, it is possible that the gamma-ray size of the 3EG J1835+5918 is considerably smaller than the physical size of a whole forming cluster. This consideration also suggests a possibility that some of gamma-ray clusters may be observed as multiple sources within $\sim \theta_{vir}$, that may be revealed by future gamma-ray missions.

One of the characteristics that the source of 3EG J1835+5918 must have is very weak radio flux that is at least two orders of magnitudes fainter than any of the securely identified EGRET blazars (Mirabal et al. 2000). The spectrum of blazars is well understood by the two components of radiation by the same population of non-thermal electrons, i.e., synchrotron radiation in radio, optical, and X-ray bands, and inverse-Compton radiation in GeV and TeV gamma-ray bands (e.g., Inoue & Takahara 1996; Kataoka et al. 1999). The ratio of luminosities

by the two processes is, as is well-known, given by the ratio of magnetic energy density to the target photon energy density, that is typically of order unity for blazars. On the other hand, this ratio is $U_B/U_{CMB} \sim 2.7 \times 10^{-3}(\xi_B/10^{-3})(M/10^{15}M_\odot)^{2/3}$ for gamma-ray clusters, by using the expression of B given in §2. Therefore, the U_B/U_{CMB} ratio is generally much smaller than the unity, and very weak radio flux compared with identified blazars can be reasonably explained.

Based on the above arguments, we suggest that the uncatalogued x-ray cluster near 3EG J1835+5918 may be a gamma-ray cluster proposed in this paper, and further observations for this cluster and the surrounding region are very important. Our model predicts that a cluster emitting a flux of $\sim 6.06 \times 10^{-7}$ photons $\text{cm}^{-2}\text{s}^{-1}$ above 100 MeV (the flux of 3EG J1835+5918, Hartman et al. 1999) at $z = 0.102$ should have a total mass of $\sim 7 \times 10^{15}M_\odot$ and $r_{vir} \sim 5$ Mpc ($\theta_{vir} \sim 0.7^\circ$).

6. CONCLUSIONS

In this paper we have proposed a new candidate of unidentified EGRET sources: gamma-ray clusters that are just dynamically forming and emit gamma-rays due to inverse-Compton scattering of CMB photons by shock-accelerated electrons. Based on the standard theory of structure formation and assuming the injection of $\sim 5\%$ of shock energy at the formation into nonthermal electrons, we have shown that a few tens of such clusters should have already been detected by EGRET, and a significant fraction of the isotropic component of unidentified EGRET sources can be accounted for. Such gamma-ray clusters are expected to be very extended; the x-ray surface brightness and surface number density of galaxies could be lower than those of ordinary clusters by a factor of ~ 200 and ~ 30 , respectively. Therefore it should have been very difficult to detect gamma-ray clusters in the past x-ray or optical surveys, and our scenario is in accord with apparent no-associations between unidentified EGRET sources and x-ray or optical clusters.

It will be of great significance to perform x-ray or optical observations to search for such loose clusters of galaxies in the regions of high-latitude unidentified EGRET sources. The future gamma-ray projects such as GLAST will also provide a direct test of our scenario. If our scenario is true, a new population of “gamma-ray clusters” will provide us in the future a new probe of dynamically evolving structures in the universe that cannot be traced by x-ray or optical clusters of galaxies.

We would like to thank S. Inoue and T. Naito for useful discussions. TT has partially been supported by the Grant-in-Aid for the Scientific Research Fund (No. 12047233) of the Ministry of Education, Science, and Culture of Japan. TK has been supported in part by the Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists.

REFERENCES

- Abramopoulos, F. & Ku, W. 1983, *ApJ*, 271, 446
 Abell, G.O., Corwin, H.G., Olowin, R.P., 1989, *ApJS*, 70, 1
 Adami, C., Mazure, A., Katgert, P., & Biviano, A. 1998, *A&A*, 336, 63
 Cen, R. & Ostriker, J.P. 1999, *ApJ*, 514, 1
 Coppi, P.S. & Aharonian, F.A. 1997, *ApJ*, 487, L9
 de Bernardis, P. et al. 2000, *Nature*, 404, 955
 Ebeling, H. et al. 1998, *MNRAS*, 301, 881
 Eke, V.R., Cole, S. & Frenk, C.S. 1996, *MNRAS*, 282, 263
 Fusco-Femiano, R. et al. 1999, *ApJ*, 513, L21
 Gehrels, N. & Michelson, P. 1999, *Astropart. Phys.* 11, 277
 Gehrels, N., Macomb, D.J., Bertsch, D.L., Thompson, D.J., & Hartman, R.C. 2000, *Nature*, 404, 363
 Hartman, R.C. et al. 1999, *ApJS*, 123, 79
 Inoue, S. & Takahara, F. 1996, *ApJ*, 463, 555
 Jones, C. & Forman, W. 1984, *ApJ*, 276, 38
 Kataoka, J. et al. 1999, *ApJ*, 514, 138
 Kitayama, T. & Suto, Y. 1996a, *MNRAS*, 280, 638
 Kitayama, T. & Suto, Y. 1996b, *ApJ*, 469, 480
 Kulsrud, R.M., Cen, R., Ostriker, J.P. & Ryu, D. 1997, *ApJ*, 480, 481
 Koyama, K. et al. 1995, *Nature*, 378, 255
 Kronberg, P. 1994, *Rep. Prog. Phys.* 57, 325
 Lacey, C. & Cole, S. 1993, *MNRAS*, 262, 627
 Loeb, A. & Waxman, E. 2000, *Nature*, 405, 156
 Mirabal, N., Halpern, J.P., Eracleous, M., & R.H. Becker 2000, to appear in *ApJ*, astro-ph/0005256
 Mücke, A. & Pohl, M. 2000, *MNRAS* 312, 177
 Mukherjee, R. et al. 1995, *ApJ*, 441, L61
 Özel, M.E. & Thompson, D.J. 1996, *ApJ*, 463, 105
 Percival, W. & Miller, L. 1999, *MNRAS*, 309, 823
 Peebles, P.J.E., 1980, *The Large-Scale Structure of the Universe*. Princeton Univ. Press, Princeton
 Press, W.H. & Schechter, P. 1974, *ApJ*, 187, 425 (PS)
 Rephaeli, Y., Gruber, D., & Blanco, P. 1999, *ApJ*, 511, L21
 Sasaki, S. 1994, *PASJ*, 46, 427
 Sarazin, C.L. 1988, *X-ray Emission from Clusters of Galaxies*. Cambridge, Cambridge Univ. Press.
 Sreekumar, P. et al. 1996, *ApJ*, 464, 628
 Sreekumar, P. et al. 1998, *ApJ*, 494, 523
 Tanimori, T. et al. 1998, *ApJ*, 497, L25
 Totani, T. 2000, *ApJ*, 536, L23